AMENDMENTS TO THE CLAIMS

The following Listing of Claims replaces all prior listings of claims in this application.

Listing of Claims:

- 1. (Currently amended) A method of killing microorganisms in aqueous systems or products based on water comprising adding a biocide to the system or product, wherein the biocide comprises from 0.001 to 5% by weight of at least one water-soluble or water-dispersible polymer comprising
 - (a) from 30 to 98 mol% of styrenesulfonic acid,
 - (b) from 2 to 40 mol% of an N-vinyllactam, N-vinylamine, or a mixture thereof, and
- (c) from 0 to 30 mol% of free-radically polymerizable monomers, wherein the mol% is based on the total molar amount of monomer units present in the polymer, and the sum of (a), (b), and (c) totals 100 mol%.
- 2. (Original) A method as claimed in claim 1, wherein all or some of the sulfonic acid groups are in salt form.
- 3. (Previously presented) A method as claimed in claim 1, wherein the products based on water are aqueous dispersions.
- 4. (Original) A method as claimed in claim 3, wherein the dispersion is electrostatically or ionically stabilized.
 - 5. (Original) A method as claimed in claim 3, wherein the dispersion is spray dried.
- 6. (Currently amended) A method as claimed in claim 1, wherein the aqueous industrial systems are refrigeration or heat exchanger circuits.

Application No. 10/529,629 Docket No.: 12810-00042-US1 Amendment dated December 12, 2008

Reply to Office Action of October 29, 2008

7. (Currently amended) A method of protecting articles by applying an antimicrobial composition comprising water or a predominantly hydrous solvent mixture and a biocide to the article and removing the water or the predominantly hydrous solvent mixture, wherein the biocide is from 0.001 to 5% by weight of at least one water-soluble or water-dispersible polymer comprising

- (a) from 30 to 98 mol% of styrenesulfonic acid,
- (b) from 2 to 40 mol% of an N-vinyllactam, N-vinylamine, or a mixture thereof and
- (c) from 0 to 30 mol% of free-radically polymerizable monomers, wherein the mol% is based on the total molar amount of all monomer units present in the polymer, and the sum of (a), (b), and (c) totals 100 mol%.
- 8. (Original) A method as claimed in claim 7, wherein the antimicrobial composition further comprises at least one binder.
- 9. (Previously presented) A method as claimed in claim 7, wherein the antimicrobial composition further comprises a crosslinker or a system of crosslinkers.
- 10. (Currently amended) A method as claimed in claim 1 wherein the water-soluble and water-dispersible polymer comprises 50 to 90 mol% styrenesalfonic styrenesulfonic acid, and 3 to 30 mol% N-vinyllactam, N-vinylamine or a mixture thereof.
- 11. (Currently amended) A method as claimed in claim 1 wherein the free-radically polymeritable polymerizable monomers contain crosslinkable groups.
- 12. (Currently amended) A method as claimed in claim 11 wherein the free-radically polymerizable monomer is acetoacetoxyethyl methacrylate.
- 13. (Currently amended) A method as claimed in claim 10 wherein the free-radically polymeritable polymerizable monomer is present from 5 to 15 mol%.

Application No. 10/529,629 Docket No.: 12810-00042-US1 Amendment dated December 12, 2008

Reply to Office Action of October 29, 2008

14. (Previously presented) A method as claimed in claim 10 wherein the at least one

water-soluble or water-dispersible polymer has a polydispersity M_w/M_n from 1.3 to less than 3.

15. (Currently amended) A method as claimed in claim 7 wherein the water-soluble

and water-dispersible polymer comprises 50 to 90 mol% styrenesalfonic styrenesalfonic acid,

and 3 to 30 mol% N-vinyllactam, N-vinylamine or a mixture thereof.

16. (Currently amended) A method as claimed in claim 7 wherein the free-radically

polymeritable polymerizable monomers contain crosslinkable groups.

17. (Currently amended) A method as claimed in claim 16 wherein the free-radically

polymeritable polymerizable monomer is acetoacetoxyethyl methacrylate.

18. (Currently amended) A method as claimed in claim 15 wherein the free-radically

polymeritable-polymerizable monomer is present from 5 to 15 mol%.

19. (Previously presented) A method as claimed in claim 15 wherein the at least one

water-soluble or water-dispersible polymer has a polydispersity M_w/M_n from 1.3 to less than 3.

4